

hep-ph/0107313

KUNS-1729

February 1, 2008

A natural solution for the μ problem with anomalous $U(1)_A$ gauge symmetry

Nobuhiro Maekawa^a

*Department of Physics, Kyoto University,
Kyoto 606-8502, Japan*

Abstract

Recently we proposed an attractive scenario of grand unified theories with anomalous $U(1)_A$ gauge symmetry, in which doublet-triplet splitting is naturally realized in $SO(10)$ unification using Dimopoulos-Wilczek mechanism and realistic quark and lepton mass matrices can be obtained in a simple way. In this paper we show that there is a mechanism in which the doublet Higgs obtains the supersymmetric mass which is proportional to the SUSY breaking parameters. This mechanism can be applied easily in the above scenario. The point is that the mass term, which is forbidden by SUSY zero mechanism, can be induced by SUSY breaking. The proportional coefficient is controlled by the anomalous $U(1)_A$ charges.

^ae-mail: maekawa@gauge.scphys.kyoto-u.ac.jp

Recently we proposed an attractive scenario of supersymmetric (SUSY) grand unified theories (GUTs) with anomalous $U(1)_A$ gauge symmetry, in which doublet-triplet splitting is naturally realized in $SO(10)$ unification using Dimopoulos-Wilczek mechanism, and realistic quark and lepton mass matrices can be obtained in a simple way [1].

In the scenario, the mass term of the Higgs field is forbidden by the holomorphy (SUSY zero mechanism). This is because the anomalous $U(1)_A$ charge of the Higgs field is taken to be negative. However, in order to give a mass to higgsino, the SUSY Higgs mass term is required. The SUSY Higgs mass μ must be of order of the weak scale, namely, the SUSY breaking scale. This is a mystery in the minimal SUSY standard model, because at a glance we have no reason that the SUSY parameter becomes the same order of the SUSY breaking parameters. This is called the μ problem [2]. In the super gravity scenario, there are several natural solutions for the μ problem [3, 4] by using non-renormalization operator in the Kähler potential (Giudice and Masiero [3]) or in the superpotential (Casas and Muñoz [4]). However, if the Higgs mass term in tree level is forbidden by some symmetry as in our scenario, these mechanisms for the μ problem do not work well, though R-symmetry can be an exceptional one. Since our model has no R-symmetry, it is important to examine other mechanisms to induce the supersymmetric Higgs mass term related with the SUSY breaking scale. There are several other attempts [5, 6, 7, 8, 9] to solve the μ problem. One of them is to introduce a light singlet which couples with the Higgs doublet [5, 7]. The vacuum expectation value (VEV) of the singlet field can become of order of the SUSY breaking scale, so suitable μ parameter is induced. In addition to the problem that the induced μ parameter is unstable under radiative correction of heavy particle and non-renormalizable terms [6], however, in our scenario, it is not easy to introduce a light singlet S with positive charge (the positive charge is required for the singlet to couple to Higgs field), because the mass term of the singlet field with positive charge is not forbidden.

In this paper, we examine a mechanism which solves the μ problem. The mechanism can be naturally applied to our scenario. The generated Higgs mass μ is proportional to the SUSY breaking parameters and the coefficient is controlled by anomalous $U(1)_A$ charges. The point is that since the Higgs mass term is forbidden by the SUSY zero mechanism, when SUSY is broken, the μ term must be induced. Note that if all the SUSY breaking parameter become zero, the μ term must vanish. Therefore the μ parameter must be proportional to the SUSY breaking parameters. Since all the coefficients are controlled by anomalous $U(1)_A$ charges, the proportional coefficient is also determined by the anomalous $U(1)_A$ charges.

Let us recall the SUSY zero mechanism. First of all, we assume that D -flatness condition of the anomalous $U(1)_A$ gauge symmetry leads to $\langle \Phi \rangle = \lambda M_P$, because $D_A = \frac{q_A}{2}(\lambda^2 M_P^2 - |\Phi|^2)$, where Φ has negative anomalous $U(1)_A$ charge $\phi = -1$ and $\lambda < 1$ (Actually we usually adopt $\lambda \sim 0.2$ for reproducing the

Cabbibo angle.). Here M_P is some gravity scale and usually taken as the reduced Planck mass, $1/\sqrt{8\pi G_N}$. In the following, we use the units in which $M_P = 1$. Then the hierarchical structure of Yukawa couplings can be obtained as

$$W_Y = \Phi^{q+u+h} Q U H \rightarrow \lambda^{q+u+h} Q U H, \quad (1)$$

if $q + u + h \leq 0$. Here q, u and h are anomalous $U(1)_A$ charges of the superfields Q, U and H ¹. The terms with negative total anomalous $U(1)_A$ charge are forbidden by the anomalous $U(1)_A$ gauge symmetry, while the terms with non-negative total charge are allowed, because the negative charge of the singlet Φ can compensate for the positive charge, as addressed above. The vanishing of the coefficients resulting from the anomalous $U(1)_A$ gauge symmetry is called SUSY zero mechanism. In the previous paper [1], this SUSY zero mechanism plays an essential role to solve the two biggest problems in grand unified theory, the doublet-triplet splitting problem and hierarchy problem of quark and lepton mass matrices. For example, the Higgs mass term μ in tree level can be forbidden by the mechanism if the anomalous $U(1)_A$ charge of the Higgs H is negative. It is obvious that the vacuum expectation values of the gauge singlet operators with positive anomalous $U(1)_A$ charges must vanish so that the SUSY zero mechanism works well. On the other hand, the gauge invariant operator with negative charge can have the VEV. The value is written

$$\langle O \rangle \sim \lambda^{-o}, \quad (2)$$

if the F -flatness condition determines the VEV [1].

Generally if SUSY is broken, the coefficients, which vanish by SUSY zero mechanism, become tiny non-zero values which are proportional to the SUSY breaking parameters. In the followings, we estimate the coefficients by anomalous $U(1)_A$ charges.

Before examining this mechanism, we try to apply the Giudice-Masiero mechanism to induce the SUSY Higgs mass μ . When SUSY is broken by the F -term F_T of a field T , a Kähler term

$$\int d\theta^4 \lambda^{|2h-t|} T^\dagger H H \quad (3)$$

induces the SUSY Higgs mass term

$$\int d\theta^2 \mu H H = \int d\theta^2 \lambda^{|2h-t|} F_T^\dagger H H. \quad (4)$$

Here H is the Higgs superfield, whose representation is, for example, **10** in the context of $SO(10)$ unification. We have to take $t = 0$ so that the gaugino mass

¹ Throughout this paper we denote all the superfields and the scalar component fields with uppercase letters and their anomalous $U(1)_A$ charges with the corresponding lowercase letters.

is obtained by $\langle F_T \rangle \sim m_{SB} M_P$, because the supersymmetric field strength has vanishing anomalous $U(1)_A$ charge. Here m_{SB} is a typical SUSY breaking scale, which is of order of the weak scale. Then the induced Higgs mass term has suppression factor $\lambda^{|2h|}$, so it is much smaller than the weak scale unless $|2h| \leq 1$ [9].

We now examine the solution for the μ problem in a simple example. The essential point of this mechanism is that the VEV shift of a heavy singlet field by SUSY breaking. In the literature [10], it is shown that the SUSY breaking terms produce the VEV shift of heavy particles of order the SUSY breaking scale in the context of super gravity scenario. If there is a heavy singlet which has vanishing VEV in SUSY limit and couples to the Higgs field, then shifting the VEV solves the μ problem. The argument is essentially the same as in Ref.[8], but they requires R-symmetry, which is not a symmetry in our scenario [1]. Below we show that such a situation is easily obtained in our scenario, namely, R-symmetry is not an essential ingredient of the mechanism.

Before examining the detail, we figure out the essence of the mechanism. We introduce the superpotential $W = \lambda^s S + \lambda^{s+z} SZ$, where S and Z are singlet fields with positive anomalous $U(1)_A$ charge s and with negative charge z , respectively ($s + z \geq 0$). Note that the single term of Z is not allowed by SUSY zero mechanism, while usual symmetry cannot forbid this term. This is an essential point of this mechanism. The SUSY vacuum is at $\langle S \rangle = 0$ and $\langle Z \rangle = \lambda^{-z}$. After SUSY is broken, these VEVs are modified. To determine the VEV shift of S , which we would like to know because the singlet S with positive charge can couple to the Higgs field with negative charge, the most important SUSY breaking term is the tadpole term of S , namely $\lambda^s M_P^2 AS$. Here A is a SUSY breaking parameter of order of the weak scale. By this tadpole term, the VEV of S appears as $\langle S \rangle = \lambda^{-s-2z} A$. If we have $\lambda^{s+2h} SH^2$, the SUSY Higgs mass is obtained as $\mu = \lambda^{2h-2z} m_{SB}$, which is proportional to the SUSY breaking parameter m_{SB} and the proportional coefficient can be of order 1 if $h \sim z$.

Let us examine the detail below. The superpotential is written

$$W = \lambda^{s+2h} SH^2 + \lambda^s S + \lambda^{s+z} SZ + \lambda^{2s} S^2, \quad (5)$$

where for simplicity, we introduce only the last term for breaking the R-symmetry. Introducing the other R-symmetry breaking terms does not change the following results drastically. The F -flatness conditions are

$$F_S = \frac{\partial W}{\partial S} = \lambda^{s+2h} H^2 + \lambda^s + \lambda^{s+z} Z + 2\lambda^{2s} S = 0, \quad (6)$$

$$F_Z = \frac{\partial W}{\partial Z} = \lambda^{s+z} S = 0, \quad (7)$$

$$F_H = \frac{\partial W}{\partial H} = \lambda^{s+2h} SH = 0. \quad (8)$$

Since $S = 0$ satisfy the two F -flatness conditions $F_Z = F_H = 0$, the VEVs of the other fields Z and H are not fixed completely by these F -flatness conditions.

Though the desired VEV is $(\langle Z \rangle, \langle H^2 \rangle) \sim (\lambda^{-z}, 0)$, the VEV of H^2 can have non-vanishing value. (Here only for simplicity, we assume that the D -flatness condition of the anomalous $U(1)_A$ gauge symmetry determine the VEV $\langle \Phi \rangle \sim \lambda$. However, in principle, the Froggatt-Nielsen field Φ is a dynamical variable, so we have to resolve the D -flatness condition in addition to the above F -flatness conditions. We will discuss this point later.)

After SUSY is broken, the SUSY breaking terms are given as

$$V_{SB} = m_S^2 |S|^2 + m_Z^2 |Z|^2 + m_H^2 |H|^2 \quad (9)$$

$$+ (\lambda^{s+2h} A_{SH^2} S H^2 + \lambda^s A_S S + \lambda^{s+z} A_{SZ} S Z + \lambda^{2s} A_{S^2} S^2 + h.c.). \quad (10)$$

Here m_X and A_Y ($X = S, Z, H$ and $Y = SH^2, S, SZ, S^2$) are SUSY breaking parameters. If we neglect the D -term contribution to the potential, the potential is obtained by

$$V = \left| \frac{\partial W}{\partial S} \right|^2 + \left| \frac{\partial W}{\partial H} \right|^2 + \left| \frac{\partial W}{\partial Z} \right|^2 + V_{SB}. \quad (11)$$

Here we assume that the Kähler potential is minimal one $K = |H|^2 + |Z|^2 + |S|^2$ for simplicity, but more general Kähler potential does not change the following result drastically unless the Kähler potential has a singularity. The stationary conditions are written

$$\frac{\partial V}{\partial S} = 2F_S^\dagger \lambda^s + \lambda^{2s+2z} S^\dagger + 4\lambda^{2s+4h} |H|^2 S^\dagger + m_S^2 S^\dagger + \lambda^{s+2h} A_{SH^2} H^2 \quad (12)$$

$$+ \lambda^{s+z} A_{SZ} Z + 2\lambda^s A_S = 0, \quad (13)$$

$$\frac{\partial V}{\partial Z} = F_S^\dagger \lambda^{s+z} + m_Z^2 Z^\dagger + \lambda^{s+z} A_{SZ} S = 0, \quad (14)$$

$$\frac{\partial V}{\partial H} = 2F_S^\dagger \lambda^{s+2h} H + 2\lambda^{2s+4h} |S|^2 H^\dagger + m_H^2 H^\dagger + 2\lambda^{s+2h} A_{SH^2} S H = 0. \quad (15)$$

The third condition (15) leads to the following two cases; a) $\langle H \rangle = 0$ and b) $\langle H \rangle \neq 0$. The second condition (14) determines the F term of the S field $F_S = -\lambda^{-s-z} m_Z^2 Z^\dagger - A_{SZ} S$, which is of order m_{SB}^2 if $\langle S \rangle \sim O(m_{SB})$. This is important to induce the correct size of B parameter. It is easily checked that with vanishing SUSY breaking parameters, the above three conditions become three F -flatness conditions $F_S = F_H = F_Z = 0$. In the following, we assume that the vacuum can be expanded as $\langle X \rangle = X_0 + X_1 + \dots$ ($X = S, Z, H$) using the SUSY breaking scale as the expansion parameter. Here X_0 represent SUSY vacua, namely, $S_0 = 0$ and $F_S(S_0, Z_0, H_0) = 0$. The first condition (13) gives

$$S_1 = -\frac{\lambda^s A_S + \lambda^{s+z} A_{SZ} Z_0}{\lambda^{2s+2z} + 4\lambda^{2s+4h} |H_0|^2}. \quad (16)$$

In the case a) ($\langle H \rangle = 0$), the relation $F_S(S_0, Z_0, H_0) = 0$ leads to $Z_0 = \lambda^{-z}$. Then from eq.(16), S_1 is given by $S_1 = \lambda^{-s-2z} A$. At the vacuum, the value of the

potential is roughly estimated as

$$V_a \sim \lambda^{-2z} |A|^2. \quad (17)$$

Let us consider the case b) ($\langle H \rangle \neq 0$). If the VEV $\langle Z \rangle \gg \lambda^{-z}$, the value of the potential $V_b \sim m_Z^2 |Z|^2 \gg V_a$. Therefore we take $Z_0 \sim O(\lambda^{-z})$. Then the equation $F_S(S_0, Z_0, H_0) = 0$ leads to $H_0 \sim O(\lambda^{-h})$. From the eq. (16),

$$S_1 = -\frac{\lambda^s A_S + \lambda^s A_{SZ}}{\lambda^{2s+2z} + 4\lambda^{2s+2h}} \sim \begin{cases} \lambda^{-s-2z} A & (z \leq h) \\ \lambda^{-s-2h} A & (z \geq h). \end{cases} \quad (18)$$

Then $2F_S^\dagger \lambda^{s+2h} + 4\lambda^{2s+4h} |S_1|^2 + m_H^2 + 2\lambda^{s+2h} A_{SH^2} S_1 = 0$ and $F_S(S_0, Z_0, H_0) = 0$ determine the H_0 and Z_0 definitely. Since all VEVs are determined, the value of the potential at the minimum can be estimated as

$$V_b \sim \begin{cases} \lambda^{-2z} |A|^2 (z \leq h) \\ \lambda^{-2h} |A|^2 (z \geq h). \end{cases} \quad (19)$$

Therefore $z > h$ leads to $V_a > V_b$, namely the desired vacuum (case a) becomes local minimum. On the other hand, if $z \leq h$, we obtain $V_a \sim V_b$. Therefore, the desired vacuum (case a) can be global minimum though it is dependent on the $O(1)$ coefficients.

Below we focus on the desired vacuum (case a) even if the vacuum is local minimum. Then the VEV of S induces the supersymmetric Higgs mass term μ as

$$\lambda^{s+2h} S H^2 \rightarrow \lambda^{s+2h} \langle S \rangle H^2 = \lambda^{2h-2z} A H^2. \quad (20)$$

It is interesting that the μ term is proportional to the SUSY breaking parameter A . The proportional coefficient is determined by the anomalous $U(1)_A$ charges as λ^{2h-2z} . When $h \sim z$, we can obtain the natural scale of the SUSY Higgs mass μ . The Higgs mixing term $B\mu$ can be obtained from the SUSY term $\lambda^{s+2h} S H^2$ and the SUSY breaking term $\lambda^{s+2h} A_{SH^2} S H^2$ as $\lambda^{s+2h} F_S \sim \lambda^{2h-2z} m_{SB}^2$ and $\lambda^{2h-2z} A^2 \sim \mu A$, respectively. Therefore the relation $B \sim m_{SB}$ is naturally obtained². This is a solution for the μ problem. Note that the condition $h \sim z$ can be satisfied because both fields H and Z have negative charges.

At a glance, requiring the condition $h \sim z$ is artificial. However, recall that even the Giudice-Masiero mechanism requires an additional condition $h \sim 0$.

In the above argument, we almost fix the VEV $\langle \Phi \rangle \sim \lambda$, which is considered to be determined by the D -flatness condition of the anomalous $U(1)_A$ gauge symmetry. Since the Froggatt-Nielsen field Φ is a dynamical variable, in principle, we have to reconsider the D -flatness condition of anomalous $U(1)_A$ gauge

² If doublet-triplet splitting is realized by fine-tuning or some accidental cancellation, the Higgs mixing $B\mu$ can become intermediated scale $m_{SB} M_X$ as discussed in Ref.[11], where M_X is the GUT scale. However, once the doublet-triplet splitting is naturally solved as in Ref.[1], such a problem disappears

symmetry to determine the VEVs. However, the result is almost the same as that discussed in my paper if $\langle \Phi \rangle \sim \lambda$ and $z < -1$. Since the VEV of Z is λ^{-z} , which is much smaller than the VEV of Φ , reconsidering D -flatness condition $\lambda^2 - |\Phi|^2 - z|Z|^2 = 0$ makes only a tiny shift in the VEV of Φ . Of course there is another possibility that the other vacuum appears, for example, $\langle \Phi \rangle = 0$ and $\langle Z \rangle \sim \lambda$ (case c). (Actually this vacuum satisfies all the F -flatness conditions (6), (7) and (8).) In such a case, the role of Φ is exchanged for that of Z . The condition $\phi = -1 > h$ leads to $V_c \sim \min(\lambda^{2h/z}|A|^2, \lambda^2 m_{SB}^2) > \lambda^{-2h}|A|^2 \sim V_b$, which means that the vacuum c is an only local minimum. Moreover, if we adopt the charges $s = -nz$ (n is a positive integer), then the F -flatness condition of S requires $\Phi^s + Z^n \sim 0$, namely, $\Phi \sim Z^{-1/z} > \lambda$, which is inconsistent with D -flatness condition. Then the vacuum c is not allowed. In any cases, the situation is not changed drastically by examining the D -flatness condition. This is consistent with the number of equations and variables. If we add one equation (D -flatness condition) and one variable (Φ), it is expected that the number of the vacua does not increase drastically.

In this paper, we have focused on the μ problem. However, the mechanism can be applied to more general case. We can apply this mechanism to induce any mass term which is forbidden by SUSY zero mechanism. Namely, we can give masses to any fields which have no mass term from the superpotential with SUSY vacua.

In summary, we have examined a solution for the μ problem. The point is that if SUSY is broken, the Higgs mass term, which is forbidden by holomorphy (the SUSY zero mechanism), must be induced. It is interesting that the proportional coefficient is determined by anomalous $U(1)_A$ charges. The result is independent on the detail of the mediation mechanism of the SUSY breaking. We only assume that all the SUSY breaking parameters are given as in eq. (10). This is a remarkable feature of this mechanism.

Note added: After almost finishing this work, we noticed a recent paper by Kitano and Okada [12], in which a solution for the μ problem is discussed. Their solution has a similar structure of our solution, but is different. Their essential point is that when SUSY is broken, the R -symmetry, which forbids the SUSY Higgs mass term in tree level, is also broken³. Then the Higgs mass term can be induced by super gravity effect. Their solution requires the R -symmetry, which gives a severe constraint to the possible interactions. Actually it cannot apply to our scenario in which R -symmetry does not exist. On the other hand, since the mechanism discussed in this paper requires only the holomorphic zero (SUSY zero) mechanism, it can apply to our scenario[1].

We would like to thank H. Nakano for pointing us to Ref.[11].

³This is the same argument as in Ref.[8], though they succeeded to omit the additional symmetry which is introduced in Ref. [8].

References

- [1] N. Maekawa, hep-ph/0104200, to be published in *Prog. Theor. Phys.*.
- [2] J. Kim and H.P. Nilles, *Phys. Lett.* **138B** (1984) 150.
- [3] G.F. Giudice and A. Masiero, *Phys. Lett.* **B206** (1988) 480.
- [4] J.A. Casas and C. Muñoz, *Phys. Lett.* **B306** (1993) 288.
- [5] H.P. Nilles, M. Srednicki and D. Wyler, *Phys. Lett.* **B120** (1983) 346; J.-P. Derendinger and C.A. Savoy, *Nucl. Phys.* **B237** (1984) 364.
- [6] H.P. Nilles, M. Srednicki and D. Wyler, *Phys. Lett.* **B124** (1983) 337; A. Lahanas, *Phys. Lett.* **B124** (1983) 341; U. Ellwanger, *Phys. Lett.* **B133** (1983) 187; J. Bagger and E. Poppitz, *Phys. Rev. Lett.* **71** (1993) 2380; J. Bagger, E. Poppitz and L. Randall, *Nucl. Phys.* **B455** (1995) 59.
- [7] C. Kolda, S. Pokorski and N. Polonsky, *Phys.Rev.Lett.* **80** (1998) 5263.
- [8] R. Hempfling, *Phys. Lett.* **B329** (1994) 222.
- [9] Y. Nir, *Phys.Lett.* **B354** (1995) 107.
- [10] L. Hall, J. Lykken and S. Weinberg, *Phys. Rev.* **D27** (1983) 2359.
- [11] Y. Kawamura, H. Murayama and M. Yamaguchi, *Phys.Rev.* **D51** (1995) 1337.
- [12] R. Kitano and N. Okada, hep-ph/0107084.